

How to use syringes and cups to feed your baby

Information for parents



Most babies are able to breast/chest feed from birth, yet some may require supplementation by syringe or cup – these methods have been linked to improved rates of eventual exclusive breast/chest feeding. Artificial formula is not recommended, as does not have the same health benefits as breast/chest feeding, but may be used if clinically indicated, or if this is your preference.

All midwives, healthcare support workers and the Lime Green Team healthcare support workers can help with skin to skin, hands-off hand expression, syringe and cup feeding, and should offer this as the first option if supplementation is needed.

Syringe feeding



- From birth to day 3, or for small amounts of colostrum (no more than 0.2ml given at a time) use a 1ml disposable purple feeding syringe; if you use your own-bought syringes please ensure that the plastic bung is removed before offering to baby to avoid choking or injury.
- First wash your hands and hand express colostrum or collect your pre-expressed milk from the ward freezer. If the amount of milk is more than 5mls, move to cup feeding to avoid risk of choking/aspiration.
- Wrap your baby in a towel and sit them upright; maintain eye contact to aid bonding and baby's brain development.
- A member of staff will first demonstrate the technique so you feel confident. Gently stroke the syringe down over your baby's top lip to encourage rooting and mouth opening.
- Place the syringe between your baby's cheek and lower gum and push the plunger gently so your baby drinks a little at a time (0.1-2ml).

Cup feeding



- Ideal from day 3 OR quantities over 5mls.
- Only put 5-10mls of milk in the cup at a time, and refill when baby has drank the milk – there is no limit to the volume given by cup during the overall feed.
- Wrap your baby in a towel. Support your baby to sit in an upright position on your lap; maintain eye contact to aid bonding and baby's brain development.
- A member of staff will first demonstrate the technique so you feel confident.
- Gently rest the rim of the cup on your baby's lower lip – allow the milk to come UP TO the edge of the cup.
- Direct the rim of the cup towards the corners of baby's lip allowing your baby to lap the milk with their tongue – **milk should never be poured into baby's mouth**.
- Leave the cup in place to allow your baby to take what they want in their own time – a cup feed should take around 30 minutes.
- Finally, wash the cup in hot soapy water and then sterilise the cup before the next use. After removal from a cold water steriliser and before use – please rinse the cup with cooled, boiled water, or sterile water to remove residue.

Responding to your baby



- Supplements should be offered to babies in a respectful and responsive manner.
- Supplementation amounts for a healthy, term (37 weeks or above) baby may differ per feed and should reflect the normal amounts of colostrum available, the size of the infant's stomach (which changes over time), the age and size of the infant.
- Based on the limited research available, suggested intakes for term healthy infants are: **Day 0** (up to 24 hours) - 2-10 ml/per feed; **Day 1** - 5-15 ml/per feed; **Day 2** - 15-30 ml/per feed; **Day 3** - 30-60ml/per feed; **Day 5** - 30-60ml/per feed - but feeding should be by infant cue to satiation.
- SFH Maternity teams aim to address queries about your feeding plan, or listen to concerns and act as required – please use your call buzzer whilst in hospital, or phone Maternity triage on 01623 676170.

Further sources of information

NHS Choices: www.nhs.uk/conditions

Our website: www.sfh-tr.nhs.uk

Patient Experience Team (PET)

PET is available to help with any of your compliments, concerns or complaints, and will ensure a prompt and efficient service.

King's Mill Hospital: 01623 672222

Newark Hospital: 01636 685692

Email: sfh-tr.PET@nhs.net

If you would like this information in an alternative format, for example large print or easy read, or if you need help with communicating with us, for example because you use British Sign Language, please let us know. You can call the Patient Experience Team on 01623 672222 or email sfh-tr.PET@nhs.net.

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you. External websites may be referred to in specific cases. Any external websites are provided for your information and convenience. We cannot accept responsibility for the information found on them. If you require a full list of references (if relevant) for this leaflet, please email sfh-tr.patientinformation@nhs.net or telephone 01623 622515, extension 6927.

To be completed by the Communications office

Leaflet code: PIL202601-03-SCFB

Created: March 2022 / Revised: January 2026 / Review Date: January 2028