



INFORMATION FOR PATIENTS

Squint surgery - children

This leaflet will explain what a squint operation entails and what to expect while your child is in hospital. It also answers some frequently asked questions.

What does the operation entail?

Your child will have a general anaesthetic (i.e. he/she will be asleep throughout the whole procedure) and the surgeon will be operating on the muscles, which move the eyes.

Everyone has six small muscles attached to the eye which move the eyes up and down and from side to side. By strengthening or weakening some of these muscles, a squint can be corrected. The surgeon does this by shortening the muscle or by moving the position of the muscle; for example if your child has a convergent squint (where the eye turns in towards the nose), the surgeon will probably strengthen the muscle that pulls the eye out and weaken the muscle that pulls the eye in.

What happens afterwards?

On arrival back to the paediatric ward from the Day Case Unit theatre your child may have a pad over the eye for a short time. The eye/s will be very red afterwards, but most of the redness does settle down within two to four weeks of the operation.

You will be given drops or cream to put in your child's eye following the operation. These help to reduce the inflammation (redness) and guard against any infection. They also ease any discomfort as your child may complain of a gritty feeling and may not want to open the eye for the first day after the operation.

How long will my child be in hospital?

Usually your child will have the operation on the same day as he/she comes in and will go home later that day. However, if the operation is not until the afternoon, your child may still be sleepy or feeling sick from the anaesthetic and may need to stay overnight. He/she will have an appointment for pre-assessment as an outpatient the week before the operation and is seen initially after the operation either before you leave the hospital on the day of the surgery or within two days of the surgery.

How long does my child need to stay off school?

The doctor will advise you on the length of time your child needs to stay off school or nursery. This is normally about two weeks. This is to reduce the risk of infection.

Will my child still need to wear his/her glasses after the operation?

Yes, your child will still need to wear glasses unless advised otherwise.

Please remember to bring the glasses to all appointments.

Is the operation successful?

Usually the operation is very successful in straightening the eyes. However, some children need more than one operation, especially if the squint is a very large one to begin with.

Are there any problems?

As with all surgery, there is a minimal anaesthetic risk. Your child's general health will be assessed prior to surgery.

Contact details

If you have any queries about your child's treatment please ring the Orthoptic Department:

King's Mill Hospital

01623 622515, extension 3365 or 6655 Monday to Friday, 9am-4pm.

Newark Hospital

01636 681681.

Useful contact

www.orthoptics.org.uk

Further sources of information

NHS Choices: www.nhs.uk/conditions
Our website: www.sfh-tr.nhs.uk

Patient Experience Team (PET)

PET is available to help with any of your compliments, concerns or complaints, and will ensure a prompt and efficient service.

King's Mill Hospital: 01623 672222 **Newark Hospital:** 01636 685692

Email: sfh-tr.PET@nhs.net

If you need this information in a different language or format, please contact the PET (as above).

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you.

External websites may be referred to in specific cases. Any external websites are provided for your information and convenience. We cannot accept responsibility for the information found on them.

If you require a full list of references for this leaflet, please email sfh-tr.patientinformation@nhs.net or telephone 01623 622515, extension 6927.

To be completed by the Communications office

Leaflet code: PIL3249(2)

Created: November 2015 / Revised: June 2017 /

Review Date: June 2019