

INFORMATION FOR PATIENTS

4-step bowel obstruction diet

The information in this advice sheet is for patients who:

- Have been diagnosed with a bowel obstruction related to cancer.
- Have a mass in the small or large intestine (bowel) and are at risk of a blockage.

Symptoms that could be a sign your bowel may be blocked include:

- Abdominal (tummy) pain.
- Vomiting.
- Feeling full quickly.
- 'Tightness' across your stomach.
- Nausea (feeling sick).
- Bowels not opened or passed wind for more than two days.
- Bloating/abdominal (tummy) swelling.
- Feeling full after drinking.

Background

Some foods are more difficult for the body to break down. When they pass through the digestive system without being broken down properly, they can risk getting caught in a narrowing of the bowel. Reintroducing oral intake following a period of obstruction is a gradual process and people progress onto different levels depending on their obstruction and symptoms. There are four stages for reintroducing diet. How long you are on each stage will be dependent on your symptoms. You may move back and forwards between stages.

The 4 steps:

Stage 1 Clear fluids	Stage 2 Liquid diet	Stage 3 Soft/ pureed diet	Stage 4 Soft low fibre diet
--------------------------------	-------------------------------	--	--

Stage 1 – clear fluids

After a complete blockage when you start to pass wind again and your symptoms settle, you will be encouraged to start sipping on clear liquids only.

Top tip for clear fluids

You may also suck on boiled sweets and mints (ensuring they are not chewed or swallowed in pieces).

Types	Options
Sweet	Water, coconut water, black tea/ black coffee, herbal tea, fruit tea, squash, smooth/ clear fruit juice, flavoured water (Oasis®, Rubicon® (still), Vimto®), still isotonic sports drinks (Lucozade®, Powerade®).
Savoury	Clear miso soup, hot cup of water with Marmite®, a stock cube or Bovril®, clear consommé soup (no bits).

Stage 2 – liquid diet

Once your bowels have started opening regularly and your symptoms from bowel obstruction have settled, you will be able to move to the next stage of liquid diet which means all types of liquids that are completely smooth and thin (no thicker than whole milk).

Top tips for a liquid diet:

- To achieve the appropriate consistency, a liquidiser/ handheld blender can be used alongside a sieve for straining.
- To add additional calories and protein to your liquid diet, use full fat milk (dairy or dairy free alternative), add skimmed milk powder, or add cream into thin milkshakes.
- Aim to follow a little and often approach
- You can also suck on boiled sweets (not chewed or swallowed in pieces), mints, ice lollies or melt in the mouth smooth chocolate or crisps for flavour.

Mealtimes	Options
Breakfast	Blended porridge with warm milk, thin milkshake, smooth/ thin smoothie.
Main meals	Smooth homemade soups e.g. chicken, oxtail, sweet potato or butternut squash, powdered soups.
Snacks	Hot chocolate, Horlicks®, Ovaltine®, milky coffees/ tea, smooth ice cream, homemade milkshakes with milk/plant-based alternatives, milkshakes such as Frijj®, Yop®, Yazoo®, thin custard, ice cream, frozen yogurt, Sorbet, jelly.

Stage 3 - smooth/puréed diet

If your bowels are opening regularly and symptoms have settled after several days on stage 2, you can move onto stage 3. All foods should be a smooth consistency with no lumps. These foods should not have to be chewed due to their consistency

Top tips for a smooth/puréed diet:

- To achieve the appropriate consistency a liquidiser/handheld blender can be used alongside a sieve for straining.
- When blending the food, a meal can look more appetising if you keep the food groups separate (you could use moulds).

- Also include cream, cheese, skimmed milk powder, full fat milk, stock, gravy, or sauce to fortify the meal with calories and keep it creamy.
- Avoid the 'roughage' on fruits and vegetables (skin or large pips) or gristly meats as these are more difficult to blend to a smooth consistency.
- Aim to eat small portions but more regularly such as 5-6 small meals across the day.

Mealtimes	Options
Breakfast	'Soggy/ blended' cereal, Ready Brek® with milk, Weetabix®, pureed/ stewed fruit, smooth scrambled eggs, milkshake, smooth smoothie.
Main meals	Puréed meats and gravy, puréed fish with sauce, puréed stew, mashed potatoes with butter, the inside only of a jacket potato, puréed egg mayonnaise, cream cheese, smooth pâté, smooth soup, avocado. (Try adding garlic or chilli-infused oil when cooking at add in flavour).
Pudding	Custard, rice pudding, semolina, tapioca, mousse, Angel Delight®, crème caramel, milk-free alternative puddings, sponge cake blended with ice cream or custard, yogurts, ice creams, biscuits dunked in a hot drink, smooth nut butters, smooth hazelnut spread.
Melt-in-the-mouth options	Wotsits®, Skips®, Quavers®, Pom Bears®, Walkers®, crackers, smooth chocolate.

Stage 4 – soft low fibre diet

If your bowels continue to open regularly and you are not having any associated symptoms, you can move onto stage 4. Having a lower-fibre diet may impact your bowel habits, please speak to your medical team about starting laxatives as appropriate.

Top tips for a soft low-fibre diet:

- Avoid all bread/bread products including crumpets, muffins, and doughnuts. This is due to the doughy texture which swells and forms a ball which could block a narrow space increasing risk of obstruction.
- Ensure food is prepared soft, moist, and sloppy, for example adding extra sauce, butter, cream and custard. Ensure it is cooked well.
- Chew food well and take your time eating.
- Choose a little and often approach to eating, for example small meals/snacks throughout the day, not overloading with a large meal all at once.
- Avoid the 'roughage' on fruits and vegetables.
- Avoid beans/pulses.
- Avoid gristly meats.
- At the start, aim to have only 1 portion of fruit and 1 portion of vegetables per day only and increase if tolerated.

Stage 4	Foods to include	Foods to avoid
Meats, fish, eggs, and others	Meat: All types of plain meat or poultry e.g. pork medallions, trimmed back bacon, chicken breast, lean lamb/beef, minced meat, chicken or turkey, sliced meats, smooth meat paste/pâté. Eggs: All types. Fish: Plain fish with bones removed e.g. fresh, frozen, or tinned. Fish paste/pâté. Ensure prawns are peeled and chewed well. Meat-free options: Tofu, soy chicken, soy mince. Beans/pulses: Soft well-cooked beans and lentils without skins, e.g. lentil soup, yellow split peas, marrowfat peas. Completely smooth puréed beans, lentils, or hummus (with no skins).	Meat: Gristly meats, fatty lower quality minced meats and ribs. Fish: Fish that contains edible bones such as whitebait, tinned mackerel. Fish skin. Beans/pulses: Pulses with coarse skins or firm beans e.g. kidney beans, baked beans, chickpeas, soybeans, haricot beans or borlotti beans.
Dairy	Milk: Milk and non-dairy alternatives. Cheese: Plain hard or soft cheeses. Yogurts: Plain/smooth yogurts, fromage frais or natural/Greek style frozen yogurt. Puddings: Plain ice cream, custard, rice pudding, semolina, tapioca, blancmange, mousse, Angel Delight®, sorbet.	Avoid added nuts, dried fruit, seeds onion or fruits with pips/skins.
Cereals, rice, bread, pasta, grains, and potatoes	Cereals: Breakfast cereals soaked well in milk to a soggy consistency such as Rice Krispies®, Coco Pops®, Ready Brek®, Cornflakes®. Grains: Soft, well-cooked grains, e.g. white rice, plain couscous, polenta, tapioca. Potatoes: All potato types soft and without skins, chewed well with additional sauce such as mashed potato, potato waffles, fried potatoes. Crackers: Crackers, crisp breads, melt on the tongue crisps, breadsticks, Cornish wafers, Ritz® crackers, Tuc® crackers.	Cereals: Any with added dried fruits or nuts e.g. muesli, granola, Fruit 'n Fibre®, Crunchy Nut Cornflakes®. Wholemeal/ wholegrain and bran cereals e.g. All Bran®, Branflakes®, Weetabix®, Special K®. Rolled oats and porridge. Grains: Barley, pearl barley, spelt, quinoa, bulgar wheat, farro, freekeh, amaranth. Bread: All bread products including doughnuts, muffins, crumpets, pitta breads, croissants, pancakes, brioche. Crackers: Wholegrain or seeded crackers, crispbreads, oat cakes, corn cakes.

Biscuits and cakes	Cakes: Madeleines, madeira cake, angel slices, mini chocolate rolls, lemon slices, apple pie softened with custard, cream, custard, ice cream. Biscuits: Biscuits dunked in hot drinks, e.g., Rich Tea, malted milk, custard creams.	Cakes: Avoid added nuts, dried fruit or fruits with pips/skins e.g. fig rolls, Garibaldi®. Hot cross buns, tea cakes, fruit scones Biscuits: Wholegrain biscuits, Hobnobs®.
Fruits to have in moderation	All peeled and without pips: Apples, apricots, bananas, mango, melon, nectarines, papaya, pears, peaches, plums, watermelon. Tinned or stewed fruit, fruit compote, fresh fruit juice (no bits).	Blackberries, blackcurrants, blueberries, cherries, clementines, coconut, cranberries (fresh/dried), dates, dried apricots, figs, grapefruit, grapes, kiwi, mandarins, oranges, passionfruit, pineapple, pomegranate, prunes, raisins, raspberries, redcurrants, rhubarb, satsumas, strawberries, sultanas.
Vegetables to have in moderation	All vegetables need to be well cooked, peeled, deseeded, stalks removed: Aubergine, avocado, broccoli, butternut squash, cabbage, carrots, cauliflower, celeriac, courgette, cucumber, daikon (Asian radish), garlic puree, chilli puree, gem squash, ginger puree, horseradish puree, lemongrass puree, marrow, onion/shallots (well-cooked as a paste), parsnips, patty pan, peppers, pumpkin, spinach, swede, tomato passata, tomatoes, turnips, smooth vegetable soups.	All skins, pips, and seeds. Tough stalks of vegetables e.g. cauliflower, broccoli. Asparagus, baby corn, bean sprouts, beetroot, bok choy, broad beans, brussels sprouts, celery, chicory, chilli (fresh), coleslaw, edamame, fennel, French beans, garlic (whole, raw and undercooked), gherkins, ginger, globe artichokes, green beans, Jerusalem artichokes, kohlrabi, leeks, lemongrass, lettuce, mangetout, mushrooms, okra, olives, onion (whole, raw or undercooked), pak choy, peas, pickled onions, radishes, rocket, runner beans, salad leaves, shallots (whole, raw and undercooked), soya beans, spinach (raw), spring greens, spring onions, sugar snap peas, sweetcorn.
Others	Lemon curd, smooth marmalade, seedless jam, honey, syrup yeast extract, Bovril®/stock cube, tomato puree, herbs and spices, pesto that doesn't contain nuts, ketchup, mustard. Smooth nut butters in moderation, e.g. almond, peanut, cashew butters, Nutella®, or chocolate spread.	Whole fruit jams/conserves/ marmalade with shreds or seeds/skins. Pickles and chutney, crunchy vegetable-based dips. Tropical or Bombay mix. Fruit and/or nut mixes, sweets with fruit and nuts e.g. peanut brittle, yogurt or chocolate covered nuts/fruit.

	Fudge, smooth white or plain chocolate, jelly.	Toffee, jelly sweets, nougat, marshmallows, popcorn. Nuts and crunchy nut butters.
--	--	---

The oncology dietitian will plan to follow up. If you have any other questions or concerns regarding this diet, then you can contact your dietitian.

Dietitian:

Telephone number: 01623 676025

Further sources of information:

NHS Choices: www.nhs.uk/conditions

Our website: www.sfh-tr.nhs.uk

Patient Experience Team (PET)

PET is available to help with any of your compliments, concerns, or complaints, and will ensure a prompt and efficient service.

King's Mill Hospital: 01623 672222

Newark Hospital: 01636 685692

Email: sfh-tr.PET@nhs.net

If you would like this information in an alternative format, for example large print or easy read, or if you need help with communicating with us, for example because you use British Sign Language, please let us know. You can call the Patient Experience Team on 01623 672222 or email sfh-tr.PET@nhs.net.

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you. External websites may be referred to in specific cases. Any external websites are provided for your information and convenience. We cannot accept responsibility for the information found on them. If you require a full list of references (if relevant) for this leaflet, please email sfh-tr.patientinformation@nhs.net or telephone 01623 622515, extension 6927.

To be completed by the Communications office
Leaflet code: PIL202609-02-BOD
Created: July 2026 / Revised: September 2026 /
Review Date: September 2028