

Further sources of information

NHS Choices: www.nhs.uk/conditions

Our website: www.sfh-tr.nhs.uk

Patient Experience Team (PET)

PET is available to help with any of your compliments, concerns or complaints, and will ensure a prompt and efficient service.

King's Mill Hospital: 01623 672222

Newark Hospital: 01636 685692

Email: sfh-tr.PET@nhs.net

If you would like this information in an alternative format, for example large print or easy read, or if you need help with communicating with us, for example because you use British Sign Language, please let us know. You can call the Patient Experience Team on 01623 672222 or email sfh-tr.PET@nhs.net.

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you. External websites may be referred to in specific cases. Any external websites are provided for your information and convenience. We cannot accept responsibility for the information found on them.


If you require a full list of references (if relevant) for this leaflet, please email sfh-tr.patientinformation@nhs.net or telephone 01623 622515, extension 6927.

To be completed by the Communications office
Leaflet code: PIL202310-03-DAPCF
Created: January 2020 / Revised: October 2023/
Review Date: October 2025

INFORMATION FOR PATIENTS

Discharge advice

Paediatric clavicle fracture



Front view

Information, advice and recommendations for the injury

Your child has fractured their clavicle (collar bone). This type of fracture is common in children.

This type of fracture heals well – the only treatment needed is a sling.

We expect the collar bone to be painful for around four to six weeks. Your child may find it more comfortable to sleep sitting upright for a few days after the injury.

The shoulder and arm can be moved out the sling as comfort allows. This will usually be about two weeks after the injury but can be sooner if comfortable.

The bump over the fracture is quite normal and is produced by healing bone. This may take up to a year to disappear. If your child is older than ten years, a bump may remain.



Information, advice and recommendations from your nurses:

- When wearing the sling, monitor your child for any skin damage or red areas. If you see any please contact us (details below).
- The sling should be removed prior to your child going to sleep.
- Encourage your child to regularly move their fingers, wrist and elbow to prevent the joints becoming stiff.
- Give your child basic pain relief as needed - visit your local pharmacy or GP if you are not sure what pain relief should be given.

Problems and further advice

A fracture liaison service is available and advice can be given over the telephone.

If there are any issues or you have any concerns/questions, we advise you seek help.

Call the fracture liaison nurses on 01623 622515, extension 4114 or 6807, between 8am and 4pm, Monday to Friday.

Outside of these hours, or for urgent advice, please go to your nearest Emergency Department.