

Further sources of information

NHS Choices: www.nhs.uk/conditions

Our website: www.sfh-tr.nhs.uk

Patient Experience Team (PET)

PET is available to help with any of your compliments, concerns or complaints, and will ensure a prompt and efficient service.

King's Mill Hospital: 01623 672222

Newark Hospital: 01636 685692

Email: sfh-tr.PET@nhs.net

If you would like this information in an alternative format, for example large print or easy read, or if you need help with communicating with us, for example because you use British Sign Language, please let us know. You can call the Patient Experience Team on 01623 672222 or email sfh-tr.PET@nhs.net.

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you. External websites may be referred to in specific cases. Any external websites are provided for your information and convenience. We cannot accept responsibility for the information found on them.

If you require a full list of references (if relevant) for this leaflet, please email sfh-tr.patientinformation@nhs.net or telephone 01623 622515, extension 6927.

To be completed by the Communications office
Leaflet code: PIL202310-01-SCI
Created: October 2023/ Review Date: October 2025

Information reproduced with permission from
Cambridge University Hospitals NHS Trust

INFORMATION FOR PATIENTS

Discharge advice

Scaphoid injury



What is a scaphoid injury?

The scaphoid is a bone in the palm of your hand. It can be injured by simply falling onto an outstretched hand. Symptoms of a scaphoid injury typically include pain and tenderness in the area just below the base of the thumb.



You may have a clinic appointment made 10-14 days after you were first seen in the Emergency Department to re-x-ray your hand and wrist to check your progress.

Whilst your injury is healing your hand and wrist will be placed in a removable splint.

Swelling and bruising at the injury site is normal. Simple, over the counter pain killers and applying ice to the area will help to reduce swelling and pain. Ice should be applied, wrapped in a tea towel, for 20 minutes three to four times a day.

You should:

- Keep the splint on at all times even at night. It can be removed for hygiene reasons.
- Keep your fingers moving while wearing the splint to prevent them becoming stiff.
- Elevate your hand in the days following your injury to help prevent swelling.
- Remove any rings or jewellery on the affected hand.

You should not:

- Take part in any physical contact sports, as guided by your consultant at your clinic appointment.

Problems and further advice

A fracture liaison service is available, and advice can be given over the telephone.

If you develop any issues or have concerns/questions we advise you seek help.

Call the fracture liaison nurses on 01623 622515, extension 4114 or 6807, between 8am and 4pm, Monday to Friday.

Outside of these hours, or for urgent advice, please go to your nearest Emergency Department.