



OBSERVATIONS AND ESCALATION POLICY FOR ADULT PATIENTS

			POLICY	
Reference	CPG-TW-Obs&Esc			
Approving Body	Documentation Group			
Date Approved	8 th June 2023			
For publication to external SFH	Positive confirmation received from the approving body th			
website	the content does not risk the safety of patients or the public:			
	YES	NO	N/A	
	X			
Issue Date	15 th June 2023		1	
Version	9.0			
Summary of Changes from Previous Version	 Update process for 3000 bleep (no longer in use) CCOT is now a 24/7 service Auditing application changed from Perfect Wart to AMaT. 			
Supersedes	v8.0, CPG-TW-CGTRUS002, issued 6 th May 2020 to Review Date June 2023 (ext ¹)			
Document Category	Clinical			
Consultation Undertaken	 CCOT Hospital out of Hours Nervecentre Ward Leaders and Matrons Resus Sepsis Nurse Specialist NEWS2 Policy Group Deteriorating Patient Group 			
Date of Completion of Equality Impact Assessment	28 th April 2023			
Date of Environmental Impact Assessment (if applicable)	28 th April 2023			
Legal and/or Accreditation Implications	Not Applicable			
Target Audience	This policy covers all staff groups who are involved in caring for adult patients receiving care within the Trust (e.g. Registered Nurses and Students, Doctors, Physiotherapists, Pharmacists, Health Care Support Workers, Operating Department Practitioners).			
Review Date	June 2024			
Sponsor (Position)	Chief Nurse			
Author (Position & Name)	 Corporate Head of Nursing – Yvonne Simpson Practice Development Matron – Jackie Simpson 			
Lead Division/ Directorate	Corporate			
Lead Specialty/ Service/ Department	Nursing – Corporat			
Position of Person able to provide Further Guidance/Information	Corporate Nursing	Team		
Associated Documents/Information		Date Associated Information was		
None		N/A		
Template control		June 2020		

Title: Observations and Escalation Policy for Adult Patients Version: 9.0; Issued: June 2023



CONTENTS

Item	Title	Page
1.0	INTRODUCTION	3
2.0	POLICY STATEMENT	3
3.0	DEFINITIONS/ ABBREVIATIONS	4-7
4.0	ROLES AND RESPONSIBILITIES	7-12
5.0	APPROVAL	12
6.0	DOCUMENT REQUIREMENTS (NARRATIVE)	13-19
6.1	Observations	13
6.2	Frequency of Observations	13
6.3	Management Plan (Parameters/ Exceptions)	15
6.4	De-escalation within Nervecentre	15
6.5	Sp02 Scale 2	16
6.6	Urine output	16
6.7	NEWS2 scores and actions	17
6.8	Absence of response to escalation	17
6.9	Critical care referral	18
6.10	End of Life/ Palliative Care	18
6.11	Escalation of an unwell/ deteriorating member of the public or patient within Out-patient Settings	18
7.0	MONITORING COMPLIANCE AND EFFECTIVENESS	20
8.0	TRAINING AND IMPLEMENTATION	21
9.0	IMPACT ASSESSMENTS	21
10.0	EVIDENCE BASE (Relevant Legislation/ National Guidance) and RELATED SFHFT DOCUMENTS	21-22
11.0	KEYWORDS	22
12.0	APPENDICES (list)	23
Appendix 1a	Escalation of Observations Flowchart for Adult In-Patients – for King's Mill Site ONLY 09:00-17:00	24
Appendix 1b	Escalation of Observations Flowchart for Adult In-Patients – For King's Mill Site ONLY 17:00-0900	25
Appendix 2	Escalation of Observations Flowchart – for Mansfield Site ONLY	26
Appendix 3	Escalation of Observations Flowchart – for Newark Site ONLY	27
Appendix 4	Escalation of Observations Flowchart – for The Emergency Department ONLY	28
Appendix 5	NEWS2 Score including Sp02 Scale 2	29
Appendix 6	Post-Operative Patient Monitoring for Adult Surgical In-Patients	30
Appendix 7	SBAR Communication Tool	31
Appendix 8	Equality Impact Assessment Form	32-33
Appendix 9	Environmental Impact Assessment Form	34

Title: Observations and Escalation Policy for Adult Patients Version: 9.0; Issued: June 2023



1.0 INTRODUCTION

- 1.1 This policy is underpinned by the principle that recognises the importance of ensuring all adult patients' vital signs are accurately and promptly recorded by appropriately trained staff. Where patients' observations are outside of normal parameters or where there are signs of physiological deterioration, staff will take appropriate action to monitor the patient more closely and seek advice and support from other members of the Multidisciplinary Team. This aims to reverse or prevent further deterioration and avoidable harm to the patient. This policy is based on guidance from the National Institute for Health and Care Excellence (NICE, 2007) Guideline number 50 Acutely III Patients in Hospital. Recognition of and response to acute illness in adults in hospital, in conjunction with the Royal College of Physicians (2017). National Early Warning Score Second Edition (NEWS2) Standardising the assessments of acute illness severity in the NHS.
- 1.2 This policy aims to support the appropriate measuring, recording and escalation of patient's physiological observations in accordance with the National Early Warning Score Second Edition (NEWS2). It aims to depict the roles and responsibilities of different staff members in regards to escalating and responding to the deteriorating patient.
- 1.3 This policy is issued and maintained by the Chief Nurse (the sponsor) on behalf of the Trust, at the issue date defined on the front sheet, which supersedes and replaces all previous versions.

2.0 POLICY STATEMENT

- 2.1 **Subject matter:** This policy is limited to Sherwood Forest Hospital Foundation Trust (SFHFT) and covers:
 - The roles and responsibilities of different staff groups.
 - The monitoring and recording of patient vital signs.
 - NEWS2 scoring, escalation and the associated graded response (see <u>Appendix 1a</u>, <u>Appendix 1b</u>, <u>Appendix 2</u>, <u>Appendix 3</u> and <u>Appendix 4</u>).
- 2.2 **Staff:** This policy covers all staff groups who are involved in caring for adult patients (e.g. Registered Nurses, Doctors, Physiotherapists, Pharmacists, Health Care Support Worker, Operating Department Practitioners, Student Nurses and Nursing Associates).
- 2.3 **Exclusions:** The following areas are exempt:
 - o Paediatric patients in designated paediatric areas refer to PEWS Escalation.
 - Maternity patients refer to the Maternity Early Warning Scoring (MEWS) Guideline.
 - Patients who are on an end of life care plan and no longer require vital signs to be monitored - refer to Last Days of Life for Adults Policy.

Title: Observations and Escalation Policy for Adult Patients
Version: 9.0; Issued: June 2023 Page 3 of 34



3.0 DEFINITIONS/ ABBREVIATIONS:

ABCDE Approach	This is the systematic approach for assessing all patients:
	Airway
	Breathing
	Circulation
	D isability
	Exposure
	Visit the Resuscitation Councils website for further information.
ACVPU	This tool is used to rapidly assess the level of consciousness where
1.011	the patient is identified as
	A - Alert
	C – New onset C onfusion or delirium
	V – responding to V erbal stimuli only
	P – responding to P ainful stimuli only or
	U – U nresponsive
	Any person scoring C, V, P or U will require further, in-depth
	assessment using the Glasgow Coma Scale (GCS). These
	parameters all score 3 on NEWS2
	(Refer to the Acute Confusion/ Delirium in Adults (including Rapid
	Tranquilisation) - Guideline for Detection and Management for
	further information on confusion/ delirium).
ART	The Acute Response Team is a team of acute care professionals
ANI	including the on call Medical Registrar, Anaesthetic Registrar and
	CCOT/HOOHP.
	The ART is available 24 hours a day ONLY at the Kings Mill site to
	respond to acute deteriorations in a patient's clinical condition.
ССОТ	The Critical Care Outreach Team is a team of experienced senior
6601	nurses with a comprehensive range of critical care skills and
	Advance Life Support training. They provide support to the
	deteriorating patient within the Kings Mill site only. The CCOT team
	are on site 24hr a day 7 days a week.
Ceiling Of Treatment	Ceiling of treatment is an individualised patient plan that defines the
Cenning Of Treatment	limits of treatment the patient will receive in the event of
	deterioration. This should be formally documented in the medical
	notes/ ReSPECT form by a registrar or above.
	Tibles/ Nest Let form by a registral of above.
Continuous patient	At SFHFT continuous observation monitoring is defined as
observation monitoring	completing a full set of patient observations and assessment every
	30 minutes as a minimum.

Title: Observations and Escalation Policy for Adult Patients
Version: 9.0; Issued: June 2023 Page **4** of **34**



CRDM	Chronic Respiratory Disease Model – This is a model within
CINDIN	Nervecentre to reduce false alerts for those patients with chronic
	respiratory diseases including Chronic Obstructive Pulmonary
	Disease (COPD).
De-toggling	Describes the process of being able to de-escalate a patient's
De-togginig	observations on Nervecentre following an appropriate Management
	Plan documented in the medical notes, e.g. by 'de-toggling' CCOT,
	a patient will not be escalated to CCOT for a review.
DNACPR	Do Not Attempt Cardio-Pulmonary Resuscitation is a decision that
BNACFR	is made and formally recorded in conjunction with the patient, to
	recommend that cardiopulmonary resuscitation (CPR) is not
	attempted should the patient suffer a cardiac arrest or die. (Refer to
	the Trust's Cardiopulmonary Resuscitation (CPR) Policy (for Adult,
	Maternity and Paediatric Patients) for further information).
End of Life Model	A model that can be set on Nervecentre that dictates no observation
LING OF LITE MICHGI	frequency and has no automatic escalations.
GCS	The Glasgow Coma Score is used to assess a patient's neurological
300	status and is a more in-depth tool than the ACVPU.
	GCS should be used in all patients who score C, V, P or U on the
	ACVPU (unless the altered level of consciousness is expected e.g.
	due to the effects of sedative medication in theatre recovery).
	The GCS is recorded on Nervecentre.
I NA	Nursing Associate
NA	Nursing Associate – This role is designed to bridge the gap between HCSW's and Registered Nurses. They are registered
NA	between HCSW's and Registered Nurses. They are registered
	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities.
New onset Confusion or	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always
	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient
New onset Confusion or	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be
New onset Confusion or	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or
New onset Confusion or	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be
New onset Confusion or	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or worsening confusion, delirium or any other altered mentation should always prompt concern about potentially serious underlying causes
New onset Confusion or	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or worsening confusion, delirium or any other altered mentation should
New onset Confusion or delirium	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or worsening confusion, delirium or any other altered mentation should always prompt concern about potentially serious underlying causes and warrants urgent clinical evaluation.
New onset Confusion or delirium	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or worsening confusion, delirium or any other altered mentation should always prompt concern about potentially serious underlying causes and warrants urgent clinical evaluation. Health Care Support Worker's support the multidisciplinary team in
New onset Confusion or delirium	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or worsening confusion, delirium or any other altered mentation should always prompt concern about potentially serious underlying causes and warrants urgent clinical evaluation. Health Care Support Worker's support the multidisciplinary team in the delivery of high quality care. They address the care needs of
New onset Confusion or delirium	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or worsening confusion, delirium or any other altered mentation should always prompt concern about potentially serious underlying causes and warrants urgent clinical evaluation. Health Care Support Worker's support the multidisciplinary team in the delivery of high quality care. They address the care needs of individual patients under the direction and supervision of the
New onset Confusion or delirium HCSW	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or worsening confusion, delirium or any other altered mentation should always prompt concern about potentially serious underlying causes and warrants urgent clinical evaluation. Health Care Support Worker's support the multidisciplinary team in the delivery of high quality care. They address the care needs of individual patients under the direction and supervision of the Registered Nurse.
New onset Confusion or delirium HCSW	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or worsening confusion, delirium or any other altered mentation should always prompt concern about potentially serious underlying causes and warrants urgent clinical evaluation. Health Care Support Worker's support the multidisciplinary team in the delivery of high quality care. They address the care needs of individual patients under the direction and supervision of the Registered Nurse. Hospital Out of Hours (HOOH) provides effective and safe clinical
New onset Confusion or delirium HCSW	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or worsening confusion, delirium or any other altered mentation should always prompt concern about potentially serious underlying causes and warrants urgent clinical evaluation. Health Care Support Worker's support the multidisciplinary team in the delivery of high quality care. They address the care needs of individual patients under the direction and supervision of the Registered Nurse. Hospital Out of Hours (HOOH) provides effective and safe clinical care outside of normal working hours (17:00-08:00) Monday —
New onset Confusion or delirium HCSW	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or worsening confusion, delirium or any other altered mentation should always prompt concern about potentially serious underlying causes and warrants urgent clinical evaluation. Health Care Support Worker's support the multidisciplinary team in the delivery of high quality care. They address the care needs of individual patients under the direction and supervision of the Registered Nurse. Hospital Out of Hours (HOOH) provides effective and safe clinical care outside of normal working hours (17:00-08:00) Monday – Friday and 24 hours Saturday, Sunday and Bank Holidays. HOOH
New onset Confusion or delirium HCSW	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or worsening confusion, delirium or any other altered mentation should always prompt concern about potentially serious underlying causes and warrants urgent clinical evaluation. Health Care Support Worker's support the multidisciplinary team in the delivery of high quality care. They address the care needs of individual patients under the direction and supervision of the Registered Nurse. Hospital Out of Hours (HOOH) provides effective and safe clinical care outside of normal working hours (17:00-08:00) Monday – Friday and 24 hours Saturday, Sunday and Bank Holidays. HOOH is co-ordinated by the Hospital Out of Hours Practitioner (HOOHP).
New onset Confusion or delirium HCSW	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or worsening confusion, delirium or any other altered mentation should always prompt concern about potentially serious underlying causes and warrants urgent clinical evaluation. Health Care Support Worker's support the multidisciplinary team in the delivery of high quality care. They address the care needs of individual patients under the direction and supervision of the Registered Nurse. Hospital Out of Hours (HOOH) provides effective and safe clinical care outside of normal working hours (17:00-08:00) Monday — Friday and 24 hours Saturday, Sunday and Bank Holidays. HOOH is co-ordinated by the Hospital Out of Hours Practitioner (HOOHP). The HOOHP is a Registered Nurse with a comprehensive range of
New onset Confusion or delirium HCSW	between HCSW's and Registered Nurses. They are registered practitioners with identified roles and responsibilities. A patient may be alert but confused or disorientated. It is not always possible to determine whether the confusion is 'new when a patient presents acutely ill. Such a presentation should always be considered to be 'new' until confirmed otherwise. New-onset or worsening confusion, delirium or any other altered mentation should always prompt concern about potentially serious underlying causes and warrants urgent clinical evaluation. Health Care Support Worker's support the multidisciplinary team in the delivery of high quality care. They address the care needs of individual patients under the direction and supervision of the Registered Nurse. Hospital Out of Hours (HOOH) provides effective and safe clinical care outside of normal working hours (17:00-08:00) Monday – Friday and 24 hours Saturday, Sunday and Bank Holidays. HOOH is co-ordinated by the Hospital Out of Hours Practitioner (HOOHP). The HOOHP is a Registered Nurse with a comprehensive range of expanded roles who acts as a senior nurse clinician with the specific

Title: Observations and Escalation Policy for Adult Patients Version: 9.0; Issued: June 2023



Managamant Dian	A Management Dian con distate a national sindividual core needs
Management Plan	A Management Plan can dictate a patient's individual care needs
	outside of NEWS2 guidance, including exceptions to physiological
	parameters, observation frequency and escalation pathway. This must
	be clearly documented in the medical notes with a clear rationale and a
	date to be reviewed by a Doctor following consultation with a Speciality
Manadata wa Wital Cinna	Registrar.
Mandatory Vital Signs	The mandatory vital signs include:
	Respiration Rate (RR)
	Oxygen (O2) Saturations (Sp02)
	Inspired O2
	Blood Pressure (BP)
	Heart Rate (HR)
	Urine
	ACVPU
	Temperature
	• Pain
	Bowels
	Where appropriate the following should also be recorded/ instigated:
	- Oxygen delivery device (refer to Oxygen Policy - Prescription,
	Administration and Monitoring of Oxygen Therapy in Adults).
	- Fluid balance monitoring
	- Blood glucose levels
Nervecentre	The Trust's electronic observation recording system.
NEWS2	National Early Warning Score second edition.
NIC	Nurse In Charge.
Parameters	This is the term used across the Trust to describe the setting of
i arameters	exceptions to individual patient observation parameters outside of
	· · · · · · · · · · · · · · · · · · ·
	NEWS2 guidance. These parameters must be clearly documented
Olivia al Tarana	within the medical notes in the form of a Management Plan.
Clinical Team	The Clinical Team is defined as the Multidisciplinary team (MDT)
	who are responsible for a patient's medical needs whilst the patient
	is in hospital.
Urgent response	An urgent response requires the appropriate clinician to review a
	patient's escalation within 30minutes.
RN	A Registered Nurse who is registered with the Nursing & Midwifery
	Council.
ReSPECT	Recommended Summary Plan for Emergency Care and Treatment.
	(For further information visit the ReSPECT Policy)
SBAR	Is a tool to help facilitate, prompt and aid appropriate
	communication:
	Situation
	Background
	Actions
	Recommendations
	(See Appendix 7)

Title: Observations and Escalation Policy for Adult Patients Version: 9.0; Issued: June 2023



Speciality Registrar	A rotational speciality training post. In some areas, suitably qualified		
	clinical fellows or Trust grade doctors may replace this role.		
SP02	Peripheral Capillary Oxygen Saturation.		
Tag	A 'Tag' is a way of highlighting key pieces of information within		
	Nervecentre including Learning Disability, Dementia, Diabetes,		
	ReSPECT, DNACPR and Falls. Setting a Tag will cause a colourful		
	box with the appropriate information to be displayed on the patient's		
	main demographic screen.		
Vocera	Vocera is a real-time hands-free voice communication system.		
Ward Based Care	Ward based care is the ceiling of treatment for patients where the		
	decision has been documented within the medical notes by a		
	registrar or above that if the patient's condition deteriorates they are		
	not for escalation to Critical Care. The patient's medical needs		
	instead will be maintained at ward level where interventions such as		
	IV fluids, antibiotics, high flow oxygen or none invasive ventilation		
	may commence.		
Ward Doctor	Responsible doctor on a designated ward during the working day.		

4.0 ROLES AND RESPONSIBILITIES

4.1 All staff under the scope of this policy and prior to implementing care are responsible for:

- 4.1.1 Ensuring that the correct lawful consent has been gained and documented in accordance with the Trusts Policy for Consent to Examination, Treatment and Care.
- 4.1.2 Ensuring a two-stage test is undertaken on any patient where capacity is in doubt. If the patient is found to lack capacity, then a 'Best Interest Checklist' must be completed and the Trusts Mental Capacity Act (MCA) Policy followed.
- 4.1.3 Ensuring they are confident/ competent in the use of Nervecentre and have accessed the appropriate training or user guide.
- 4.1.4 Ensuring that the patient's identification data is accurate including Name/DOB/NHS number/D Number/Location and responsible Consultant) in line with the Policy and Procedure for the Positive Identification of Patients.
- 4.1.5 Ensuring that they are aware of what to do in the event of Nervecentre downtime (see <u>Nervecentre Business Continuity Plans</u> on the Trust Intranet Page for further information).

Title: Observations and Escalation Policy for Adult Patients
Version: 9.0; Issued: June 2023 Page **7** of **34**



4.2 The Registered Nurse is responsible for:

- 4.2.1 Ensuring that mandatory vital signs are recorded accurately on Nervecentre or in line with the area, i.e. ED, ICU and Recovery record on paper documentation.
- 4.2.2 If the recording of observations is not possible (for example the patient is off the ward/patient refusal) the Registered Nurse is responsible for recording why the observations have not been performed on Nervecentre or in the Nursing Evaluation.
- 4.2.3 Ensuring that observations are completed in a timely manner.
- 4.2.4 Ensuring that Nervecentre electronic handover is completed and up to date including any associate 'Tags' (e.g. ReSPECT and DNACPR). This helps responders to effectively triage deteriorating patients.
- 4.2.5 Ensuring that observations are recorded and escalated in line with either; this policy, an individual Management Plan within the medical notes or clinical judgement (see Appendix 1a, Appendix 2, Appendix 3 and Appendix 4 for specific escalation flow charts).
- 4.2.6 The Registered Nurse can delegate the recording of observations to a suitably trained and competent Nursing Associate/ Student Nurse or HCSW. However, the Registered Nurse remains responsible for ensuring that observations are recorded accurately and where required are escalated or de-toggled according to the patient's individual Management Plan. If the NEWS2 score is found to be elevated (3 in 1 parameter or an aggregated score of 5,6,7 or above) the Registered Nurse must personally review the patient.
- 4.2.7 If a patient's clinical condition warrants an urgent medical review and either the appropriate staff are not present on the ward or no clear Management Plan is in place, the Registered Nurse should immediately call the ART on ext. 2222. An ART call should be made for a patient where the NEWS2 score has increased to a 7 or more or if there are significant clinical concerns.
- 4.2.8 All staff should use this policy and Nervecentre in conjunction with their clinical judgement. If at any point you are concerned, have a question or need further support contact CCOT, HOOHP or a doctor directly via switch or Vocera.

Title: Observations and Escalation Policy for Adult Patients

Version: 9.0; Issued: June 2023 Page 8 of 34



4.3 The HCSW/Student Nurse/ Nursing Associate is responsible for:

- 4.3.1 Ensuring that they are competent/ confident in the undertaking and recording of observations and that they have completed all the associate competency packages.
- 4.3.2 Immediately informing the Registered Nurse if a patient has a raised NEWS2 score (3 in 1 parameter or an aggregated score of 5,6,7 or above) or if they or a relative are concerned about a patient.
- 4.3.3 Ensuring that observations are completed in a timely manner.

4.4 The Clinical Team is responsible for:

- 4.4.1 The overall management of the patient during working hours. Where relevant and appropriate, they are ultimately responsible for seeking a review by other specialists.
- 4.4.2 Clearly documenting and conveying the patient's on-going Management Plan to the ward staff. When documenting a Management Plan the following should be considered: triggers for escalation, observation frequency, ceiling of treatment, ReSPECT and resuscitation status.
- 4.4.3 In line with this policy, the expectation is that where the doctor reviewing the patient has concerns or is unable to resuscitate and stabilise the patient they seek urgent help from a senior colleague.
- 4.4.4 When an urgent senior review is required (due to clinical concerns or a new NEWS2 of 7 or above) and the senior clinician is not immediately available then an ART call would be appropriate (See ART section 4.8). Any patient who is seen by the ART must be reviewed by the Parent Team within 24hrs of the ART call.
- 4.4.5 In order for a rapid response to deteriorating patients, any patient outlying in a different speciality should be reviewed by the most immediately accessible doctor to prevent a delay in patient care.
- 4.4.6 If a patient has exceptions to NEWS2 parameters that should be followed, these should be clearly documented in the medical notes in the form of a Management Plan following consultation with the Speciality Registrar. Management Plans must state the specific individual patient parameters that the nurses should follow and who the Nurse should involve in the patient's escalation pathway.

Title: Observations and Escalation Policy for Adult Patients Version: 9.0; Issued: June 2023 Page 9 of 34



4.5 On Call Teams are responsible for:

- 4.5.1 Out of hours the care of the patient reverts to the On Call Speciality Team. The On Call Teams assume the responsibilities as described above for the Parent Team.
- 4.5.2 Ensuring that they make themselves available to the HOOHP, log into Nervecentre and respond appropriately (be aware that this may include seeing patients within a different speciality depending on work load).
- 4.5.3 If the clinician is likely to be unavailable to HOOH due to other commitments (e.g. emergency theatre) then they should ensure the HOOHP is made aware (via Vocera) prior to commencing this commitment.
- 4.5.4 For additional support (between 08:00-23:30) contact CCOT.

4.6 The Critical Care Outreach Team (CCOT) is responsible for:

- 4.6.1 Providing specialist advice, support and treatment to the deteriorating patient (within the King's Mill site only) 24hrs a day seven days a week, including weekends and bank holidays.
- 4.6.2 Any patient that deteriorates and has a NEWS2 of 5 or 6 should be escalated to CCOT via Nervecentre. If a patient has a NEWS2 of 7 or above they should be escalated to CCOT directly (via Vocera or bleep.888) and via the Nervecentre device (see Appendix 1a and Appendix 1b). Using clinical judgement if any concerns contact CCOT directly.
- 4.6.3 CCOT can be contacted at any time during their working hours for advice or support regardless of the patients NEWS2 score.
- 4.6.4 Any phone calls to CCOT must follow a structured SBAR referral (see appendix 7). Dependent on the clinical situation CCOT will then; provide phone advice, undertake a clinical review or suggest an ART call.
- 4.6.5 When performing a clinical review on a deteriorating patient CCOT will:
 - Perform a thorough review of the patient's history.
 - Perform a thorough physical examination using an ABCDE or system based approach.
 - Document the examination, outcome and advice within the medical notes.
 - Where appropriate, recommend that a doctor reviews the patient's observation parameters, frequency of observations, ceiling of treatment, ReSPECT and resuscitation status in the form of a clearly documented Management Plan within the medical notes.

Title: Observations and Escalation Policy for Adult Patients Version: 9.0; Issued: June 2023



4.7 Hospital Out of Hours Practitioner (HOOHP) is responsible for:

- 4.7.1 The triaging and prioritisation of all requests for assistance, treatment or care raised by ward areas on the HOOH electronic system (17:00-08:00, Monday to Friday and 24 hours Saturday, Sunday and Bank Holidays) and the delegation of these tasks to the most appropriate team member via Nervecentre.
- 4.7.2 Any patient that deteriorates and has a NEWS2 of 5 or 6 should be escalated to HOOHP via Nervecentre. If a patient has a NEWS2 of 7 or above they should be escalated to HOOHP directly (via Vocera or bleep.620) and via the Nervecentre device (see Appendix 1a and Appendix 1b).
- 4.7.3 Any phone calls to the HOOHP must follow a structured SBAR referral (see appendix 7). Dependent on the clinical situation HOOHP will then; provide phone advice, undertake a clinical review or suggest an ART call.
- 4.7.4 Attending all ART calls.
- 4.7.5 Mentoring, supporting and educating nursing staff and junior doctors.
- 4.7.6 The HOOHP will be the designated Advanced Life Support (ALS) qualified senior nurse on the Resuscitation Team.
- 4.7.7 The HOOHP can be contacted at any time during their working hours for advice or support regardless of the patients NEWS2 score.
- 4.7.8 See the Hospital Out of Hours Policy for further information.

4.8 The Acute Response Team (ART) is responsible for:

- 4.8.1 Reviewing the patient's clinical condition and documenting a clear Management Plan in the medical notes. This may include, but not exclusively, a set of clearly documented clinical interventions and on-going Management Plans, review of treatment of care and resuscitation status.
- 4.8.2 During the daytime (08:00-17:00) any patient who deteriorates requiring an ART call must be reviewed/ stabilised and then discussed with the responsible Medical Team and Parent Consultant. The ART must discuss the patient with the relevant Speciality Registrar who should contact the on call Consultant as required. All patients requiring an ART call must be reviewed by the Parent Team within 24hrs.

Title: Observations and Escalation Policy for Adult Patients
Version: 9.0; Issued: June 2023 Page 11 of 34



- 4.8.3 Where the Parent Consultant or Speciality Registrar or above is available or already reviewing the patient or if a Management Plan is in place within the medical notes then an ART call is not mandated.
- 4.8.4 The responsibility for the on-going medical management of the patient lies with the Parent Team once initial stabilisation has been achieved.

4.9 The Cardiac Arrest Team is responsible for:

- 4.9.1 Reviewing and assessing the patient's clinical condition following the Advanced Life Support (ALS) algorithm and providing a clear Management Plan that must be documented within the medical notes. This may include, but not exclusively, a set of clearly documented clinical interventions and on-going treatment plans, review of ceilings of treatment, ReSPECT and resuscitation status.
- 4.9.2 During the daytime (08:00-17:00) any patient who deteriorates requiring a cardiac arrest call must be reviewed/ stabilised and then discussed with the responsible Medical Team and discussed with the Parent Consultant. Out of hours the Cardiac Arrest Team must discuss the patient with the relevant Speciality Registrar who should contact the on call consultant as required.
- 4.9.3 The responsibility for the on-going medical management of the patient lies with the Parent Team once initial stabilisation has been achieved.

5.0 APPROVAL

Following appropriate consultation this policy has been approved by the Trust's Documentation Group.

Title: Observations and Escalation Policy for Adult Patients
Version: 9.0; Issued: June 2023 Page 12 of 34



6.0 DOCUMENT REQUIREMENTS (NARRATIVE)

6.1 Observations

- 6.1.1 The National Early Warning Score (NEWS2) is the physiological track and trigger score currently in use to identify deterioration in adult patients across the Trust.
- 6.1.2 On arrival to the Trust, all patients must have their vital signs performed and a NEWS2 score calculated and recorded.

6.2 Frequency of Observations

- 6.2.1 For the first 24 hours of admission patients must have a minimum of 4 hourly observations. Following that, if the NEWS2 score is 0, the minimum frequency of observations is 12 hourly at the Kings Mill Site and a minimum of once every 24 hours at Newark and Mansfield. If the NEWS2 score is elevated, then the observation frequency must be increased accordingly. See Appendix 1a, Appendix 1b, Appendix 2, Appendix 3 and Appendix 4 for further guidance on the frequency of observations when the NEWS2 score is above 0. For areas utilising Nervecentre the observation frequency will automatically be dictated in accordance with the appendix flow charts.
- 6.2.2 It is important to note that the patient's specific condition, intervention and care needs may dictate the frequency of observations outside of this policies guidance. Therefore it is essential that local/ specific policies, procedures and care plans are utilised in conjunction with this policy. This includes (but not exclusively):
 - Acute Pancreatitis in Adult Inpatients initial management Guidelines
 - **Head Injury Policy**
 - Post-operative Observations (see appendix 6 of this policy).
 - Patient Controlled Epidural Analgesia (PCEA) in Adults and the associated care plan.
 - Policy for Fentanyl Patient Controlled Analgesia Adult Patients and the associated care plan
 - Policy for Morphine Intravenous Patient Controlled Analgesia Adult Patients and the associated care plan
 - Day Case Unit Specific Nursing Care Pathways (Available on the Royal Marsden, click here)
- 6.2.3 To ensure patient safety when commencing or titrating medication, it is important to consult the BNF and the UCL Injectable Guidelines ('blue book') as these may dictate the frequency of observations or the monitoring equipment required (e.g. Cardiac Monitoring) outside of this policies guidance.

Title: Observations and Escalation Policy for Adult Patients Version: 9.0; Issued: June 2023



- 6.2.4 If staff have clinical concerns, irrespective of the NEWS2 score the frequency of observations should be increased as appropriate to the clinical concern. The increased frequency should be either manually set on Nervecentre or documented on the front of the paper observation chart by the Registered Nurse.
- 6.2.5 If the patient's Management Plan within the medical notes dictates the frequency of observations to be outside of NEWS2 and Nervecentre guidance (less frequently), observations on Nervecentre should be 'skipped' in accordance with the Management Plan and the option 'Agreed plan documented in notes' selected. If paper charts are in place, the frequency should be stated on the front of the observation chart and the reasoning explained within the nursing evaluation. The Registered Nurse is encouraged to use their professional judgement and perform more regular observations if they feel the patient's condition warrants.
- 6.2.6 If a patient is recovering from an acute deterioration, be aware that they are more susceptible to further deterioration. A high level of vigilance is encouraged to monitor the patient and it may be appropriate to continue more frequent observation outside of NEWS2/ Nervecentre guidance.
- 6.2.7 Only patients who are expected to die and where routine observations (set out in NEWS2) are not required as part of their care should be excluded from having observations recorded. As part of dying, it is not unreasonable for staff to perform an observation if it helps with symptom control and the patient's management, e.g. a temperature may be taken so an anti-pyretic can be administered to maintain comfort or oxygen saturation may be taken to guide staff. This is in line with the Trusts <u>Last Days of Life for Adults Policy</u>. You should consider if the patient is suitable for the end of life model on Nervecentre (see 6.9 for further information on End of Life).
- 6.2.8 Patients with a DNACPR but still for active treatment must still have their observations performed and be escalated as per this policy.
- 6.2.9 Patients who are receiving restrictive interventions and/or manual restraint e.g. mittens, must have their mandatory vital signs monitored before, during and after restraint in line with the Trust's <u>Policy for The Use of Restrictive Practice for Adult Patients</u>.
- 6.2.10 Patients who have been deemed to have capacity may in rare circumstances refuse for observations to be undertaken. In this circumstance, ensure the patient understands the importance and rationale for undertaking physiological observations. If the patient has capacity and still refuses, inform the Nurse in Charge and ensure that you clearly document the refusal on Nervecentre and/or in the nursing evaluation pages. If the patient does not have capacity,

Mental Capacity Act (MCA) Policy.

Title: Observations and Escalation Policy for Adult Patients
Version: 9.0; Issued: June 2023 Page **14** of **34**



6.3 Management Plan (Parameters/ Exceptions):

- 6.3.1 In some patients with multiple medical conditions and/ or chronic conditions the standard observation parameters may vary outside of NEWS2 guidance.
- 6.3.2 Any exceptions to individual patient parameters must be documented in the medical notes in the form of a Management Plan. This must be documented by a doctor following the consultation of a Speciality Registrar or above and verbally communicated to the Registered Nurse caring for the patient. A clear rationale and a date to be reviewed/ reconsidered must also be specified.
- 6.3.3 The newly defined observation parameters dictated within the Management Plan can be accepted as within normal range for the patient and therefore be considered to score a 0 for that parameter. If observations deteriorate outside of the specified ranges within the Management Plan then the score should reflect the standard NEWS2 guidance and be escalated in accordance with this policy. For example, if a Management Plan stated that a BP of 91-98 systolic was acceptable for the patient, the BP would score 0 and not require escalating. If the BP then dropped to 90/60, the usual NEWS2 score of 3 would apply as the BP is no longer within the parameters specified within the Management Plan and the patient should be escalated in line with this policy.
- 6.3.4 In order to follow the specified Management Plan within the medical notes, observations on Nervecentre may need to be de-toggled (see section 6.4 for further information) or skipped to ensure the correct escalation pathway and frequency of observations are followed.
- 6.3.5 Any patients with specific parameters should be highlighted during the ward handover processes and as part of Board Rounds.

6.4 **De-escalation within Nervecentre:**

- 6.4.1 In order to follow a specific Management Plan that has been documented in the medical notes, patient's observations can be de-toggled and therefore deescalated when clinically indicated (see section 6.3 for further information on Management Plans).
- 6.4.2 The De-escalation of observations in accordance with the patients Management Plan within Nervecentre is completed by de-toggling a specific responder within the escalations page of Nervecentre (the page after inputting observations). The process of de-toggling prevents unnecessary automatic escalations to responders. See the Nervecentre De-escalating Observations user guide for further support. (See EOLC/Palliative care 6.10)

Title: Observations and Escalation Policy for Adult Patients
Version: 9.0; Issued: June 2023 Page **15** of **34**



6.4.3 Only a Registered Nurse is able to de-toggle within Nervecentre. If a HCSW/ Student Nurse/ Nursing Associate is seeking authorisation for observations the Registered Nurse <u>must</u> de-toggle the observations if required before submitting the patient's observations.

6.5 **Sp02 Scale 2**

- 6.5.1 The Sp02 Scale 2 within the NEWS2 has been introduced to improve the ability to identify sick patients, and to reduce false alerts for those patients with chronic respiratory diseases including Chronic Obstructive Pulmonary Disease (COPD), who may normally have lower oxygen saturations. There is good evidence that those with COPD are at risk of oxygen toxicity, and that controlled oxygen (to target saturations of 88-92%) is associated with less morbidity and mortality in exacerbations than high flow oxygen without target saturations.
- 6.5.2 For those patients with or at risk of Hypercapnic Respiratory Failure, an appropriate clinician should decide if target saturations of 88-92% are appropriate. If appropriate, this target should be clearly documented within the patient's medical notes with an accompanying explanation of the reasoning for utilising this model. The patient can then be placed on the Chronic Respiratory Disease Model (CRDM) within Nervecentre.
- 6.5.3 The decision to use the CRDM within Nervecentre must be made by a competent clinical decision maker who is either a Registrar or above. A HOOHP or CCOT member can place the patient on the Nervecentre CRDM if a decision to use this model is clearly documented in the medical notes.
- 6.5.4 Staff should be aware that the NEWS 2 SpO2 Scale 2 (%) changes the associated scoring for the patient's oxygen saturations e.g. Sp02 of 88-92% will not trigger. See appendix 5 for further information regarding the associated scoring for Sp02 scale 2.
- 6.5.5 If the patient is not at risk of Hypercapnic Respiratory Failure the regular NEWS SpO2 scale 1 should be utilised.

6.6 **Urine output**

- 6.6.1 Urine output is not included in the NEWS2 score but is a mandatory observation.
- 6.6.2 Patients with a NEWS2 of 5 or higher must have a strict fluid balance maintained including all input and output. Catheterisation should be based on an individual assessment of the patient's condition and needs.

Title: Observations and Escalation Policy for Adult Patients
Version: 9.0; Issued: June 2023 Page **16** of **34**



6.6.3 Any urine output less than 0.5ml/kg/hr for longer than 2 hours (if catheterised) or within 6 hours (if not catheterised) must be escalated to the ward doctor/ CCOT or HOOH unless a clear Management Plan is in place to address this. This may be a sign of significant underlying illness e.g. sepsis.

6.7 NEWS2 scores and actions

- 6.7.1 See flowcharts <u>Appendix 1a</u> / <u>Appendix 1b</u> King's Mill Hospital, <u>Appendix 2</u> Mansfield Community Hospital, <u>Appendix 3</u> Newark Hospital, <u>Appendix 4</u> Emergency Department to understand the required actions based on the NEWS2 score.
- 6.7.2 This policy describes the immediate actions associated with an acute deterioration in a patient's clinical condition.
- 6.7.3 When a clinical review occurs, due to physiological deterioration, this should include:
 - A thorough review of the history and physical examination using an ABCDE approach or system based approach.
 - Formulation of a clear medical plan with appropriate investigations and interventions.
 - Review of ceilings of treatment, ReSPECT and resuscitation status.
 - Evidence of discussion/ involvement of senior colleagues where review occurs by junior staff.
 - Clearly documented and communicated Management Plans for ward staff to follow in case of further deterioration or failure to respond to intervention.

6.8 Absence of response to an escalation

- 6.8.1 In the absence of a response (review or telephone advice as appropriate) from the Medical Team/ HOOH or CCOT within 30mins of the initial escalation then the relevant individuals should be re-contacted by the ward nurse via bleep or Vocera for CCOT (888)/ HOOH(620), Switch for Doctor) and the patient should be escalated to a more senior doctor. If a patient's NEWS2 score has deteriorated further since the initial escalation, then the next level of escalation should be based on the new NEWS2 score.
- 6.8.2 If there is no response from a responder and you are immediately concerned then immediately call the ART on ext.2222.
- 6.8.3 Any failure to respond to an escalation should be documented in the nursing evaluation and a Datix completed using one of the following categories deemed most appropriate:

Title: Observations and Escalation Policy for Adult Patients
Version: 9.0; Issued: June 2023 Page **17** of **34**



- 1. Delay due to abnormal observations not escalated.
- 2. Failure to recognise deteriorating patient.
- 3. Actions—observations not acted upon.
- 4. Failure to record observations.
- 5. Failure to respond.

6.9 Critical Care Referral

- 6.9.1 Where indicated and appropriate a direct referral to the Critical Care Medical Team (bleep 269) should be made by the attending team.
- 6.9.2 An ART call or a review by a CCOT member does not automatically mean the patient has been referred to the Critical Care Medical Team. It is the responsibility of the Parent Team to directly refer the patient to the Critical Care Medical Team.

6.10 End of Life/Palliative Care

- 6.10.1 For many patients the intended purpose of this policy is to ensure there is adequate monitoring and escalation to provide active treatment/ care, but for other patients further treatment and care may be ineffective and often there has been a decision to offer best supportive and palliative care. (See de-escalation 6.4)
- 6.10.2 Where best supportive and palliative care is being offered, the monitoring of patients must change and be individualised to meet the patient's specific palliative needs and their priorities of care.
- 6.10.3 For patients in the last days of life, medical and nursing observations may not be required. This is supported by the Trusts clinical guidance referred to as the Red Bundle (see Last Days of Life for Adults Policy).
- 6.10.4 Where appropriate, a ReSPECT form should be completed and a ReSPECT Tag added to Nervecentre. Follow the <u>ReSPECT (Recommended Summary Plan for Emergency Care and Treatment) Policy</u> for further information.

6.11 Escalation of an unwell/ deteriorating member of the public or patient within Outpatient Settings

6.11.1 If you are immediately concerned about an individual or the individual is unresponsive dial 2222 for support. If an individual is found outside of the hospital buildings, summon help locally; send someone inside to dial 2222, consider dialling 999 if ambulance support is required for transfer. When calling 2222, a request for an ART call or cardiac arrest needs to be specified to the operator.

Title: Observations and Escalation Policy for Adult Patients Version: 9.0; Issued: June 2023



- 6.11.2 Click here to see the <u>Location of Resuscitation Trollies and the Areas they</u> Serve.
- 6.11.3 For the KTC, Clinic areas, Plaster Room and Main Entrance follow the <u>KTC</u>
 Medical assistance Protocol
 If the individual is not within any of those areas follow the Medical Emergency Site Protocol.
- 6.11.4 If an individual attends an outpatient's appointment and is found to be unwell but assessed to not be in immediate danger, the patient should have a full set of observations documented on a paper observation chart. The patient should then be transferred to the Emergency Department with a staff member for further assessment and treatment.

Title: Observations and Escalation Policy for Adult Patients
Version: 9.0; Issued: June 2023 Page **19** of **34**



7.0 MONITORING COMPLIANCE AND EFFECTIVENESS

- 7.1 Compliance with this policy across the Trust will be monitored by the Critical Care Outreach Team, led by the Matron for Critical Care on a monthly basis via the Trust's electronic observation monitoring system (Nervecentre) and monthly via the Trust's nursing metrics audit. This data will be reviewed in respect of acuity and dependency scores (Association of UK University hospitals acuity and dependency audit/ Safer Nursing Care Tool) and CCOT daily activity.
- **7.2** Additional on-going monitoring in areas where poor compliance is identified will be developed with individual ward teams with support from the Head of Nursing and Matron for the respective area.

Minimum Requirement	Responsible Individual	Process for Monitoring	Frequency of	Responsible Individual or
to be Monitored (WHAT – element of compliance or effectiveness within the document will be monitored)	(WHO – is going to monitor this element)	e.g. Audit (HOW – will this element be monitored (method used))	Monitoring (WHEN – will this element be monitored (frequency/ how often))	Committee/ Group for Review of Results (WHERE – Which individual/ committee or group will this be reported to, in what format (e.g. verbal, formal report etc.) and by who)
Observations	Ward leaders and Matrons	AMAT	Monthly	Ward Assurance Meeting
Observations and escalations	Critical Care Matron & CCOT	Descriptive stats from CCOT & Nervecentre data	Monthly	Deteriorating Patient Group

Title: Observations and Escalation Policy for Adult Patients



8.0 TRAINING AND IMPLEMENTATION

The contents of this policy will be integrated into training programmes already provided across the Trust. Training includes:

- Acute Illness Management (AIMS) multidisciplinary course (monitored by the Resuscitation Team).
- Mandatory Trust training Core Skills for Health Care Support Workers and Essential Skills for Registered Nurse/Registered Midwife/ Registered Mental Health Nurse (monitored by Training and Development Department and Ward and Department Leaders).
- Trust Induction training for all staff (monitored by Training and Development Department).
- Trust pain management study days (monitored by Pain Team).
- Nervecentre training Provided at Trust Induction, by accessing Intranet User guides or by contacting Nervecentre directly via switch/ Vocera.

9.0 IMPACT ASSESSMENTS

- 9.1 This document has been subject to an Equality Impact Assessment, see completed form at Appendix 8.
- 9.2 This document has been subject to an Environmental Impact Assessment, see completed form at Appendix 9.

10.0 EVIDENCE BASE (Relevant Legislation/ National Guidance) AND RELATED SFHFT DOCUMENTS

Evidence Base:

This policy has been developed with reference to the following guidance:

- British Medical Journal (BMJ, 2018). Comparison of Early Warning Scores in patients with COPD Exacerbation: DECAF and NEWS Score. Available at: https://thorax.bmj.com/content/74/10/941 [Accessed 10th March 2020].
- British Medical Journal (BMJ, 2010). Effects of High Flow Oxygen on Mortality in Chronic Restrictive Pulmonary Disease patients in pre-hospital settings. Available at: https://www.bmj.com/content/341/bmj.c5462 [Accessed 10th March 2020].
- Department of Health (DH, 2009). Competences for recognising and responding to the acutely ill patient in hospital. Gateway reference 11275.
- National Institute for Health and Care Excellence (NICE, 2007). Acutely III Patients in Hospital. Recognition of and response to acute illness in adults in hospital. Clinical Guideline No 50. Available at https://www.nice.org.uk/guidance/cg50 [last accessed 29 Aug 2019].
- National Institute for Health and Care Excellence (NICE, 2019). Acute Kidney Injury: Prevention, Detection and management. Available at https://www.nice.org.uk/guidance/ng148/chapter/Recommendations [Last accessed 4 March 2020].

Title: Observations and Escalation Policy for Adult Patients Version: 9.0; Issued: June 2023



- NHS Improvements (2018). SBAR Communication Tool Situation, Background, Assessment, Recommendation. Available at https://improvement.nhs.uk/resources/sbar-communication-tool/ [Last accessed 29 Aug 2019].
- Resuscitation Council (2018). The ABCDE Approach, Underlying Principles. Available at https://www.resus.org.uk/resuscitation-guidelines/abcde-approach/ [last accessed 29 Aug 2019].
- Royal College of Physicians (RCP, 2017). National Early Warning Score Second Edition (NEWS2). Standardising the assessment of acute-illness severity in the NHS. Available at https://www.rcplondon.ac.uk/projects/outputs/national-early-warning-score-news-2 [Last accessed 29 Aug 2019].

Related SFHFT Documents:

- Acute Confusion/ Delirium in Adults (including Rapid Tranquilisation) Guideline for **Detection and Management.**
- Acute Kidney Injury (AKI) Guidelines for Adults: Early Identification and Management
- Acute Pancreatitis in Adult Inpatients initial management Guidelines
- Cardiopulmonary Resuscitation (CPR) Policy (for Adult, Maternity and Paediatric Patients)
- Conscious Sedation Guideline (Adult Patients)
- Consent to Examination, Treatment and Care Policy
- **Deprivation of Liberty Safeguarding Policy**
- **Head Injury Policy**
- **Hospital Out of Hours Policy**
- Last Days of Life for Adult Policy
- Maternity Early Warning Scoring (MEWS) Guideline
- Mental Capacity Act (MCA) Policy
- **Nervecentre Business Continuity Plans**
- Nursing Care Pathway for Patients Undergoing General Surgery Procedures
- Oxygen Policy Prescription, Administration and Monitoring of Oxygen Therapy in **Adults**
- Patient Controlled Epidural Analgesia (PCEA) in Adults (non-obstetric) Policy
- PCA Policy Administration of Morphine, Fentanyl or Oxycodone via an Intravenous (IV) Patient Controlled Analgesia (PCA) System in Adults Policy
- Positive Identification of Patients Policy
- ReSPECT (Recommended Summary Plan for Emergency Care and Treatment) **Policy**
- Restraint and Restrictive Practices for Adult Patients Policy
- SEPSIS Guideline Recognition, Diagnosis and Early Management

11.0 KEYWORDS

Track and Trigger Score; Vital Signs; NEWS; NEWS2; National Early Warning Score; Ward Based Care; post-operative patient monitoring; Observation; Escalations; Deterioration.

Version: 9.0; Issued: June 2023 Page 22 of 34



12.0 APPENDICES

Appendix 1a	Escalation of Observations Flowchart for Adult In-Patients – for King's Mill
	Site ONLY <i>09:00-17:00.</i>
Appendix 1b	Escalation of Observations Flowchart for Adult In-Patients – for King's Mill
	Site ONLY 17:00-09:00.
Appendix 2	Escalation of Observations Flowchart – for Mansfield Site ONLY.
Appendix 3	Escalation of Observations Flowchart – for Newark Site ONLY.
Appendix 4	Escalation of Observation Flowchart – For The Emergency Department ONLY.
Appendix 5	NEWS2 Score.
Appendix 6	Post-Operative Patient Monitoring for Adult Surgical In-Patients.
Appendix 7	SBAR Communication Tool.
Appendix 8	Equality Impact Assessment Form.
Appendix 9	Environmental Impact Assessment Form

09.00-17.00 Mon-Fri Excluding bank holidays



Escalation of Observation Flowcharts – For King's Mill Site ONLY

High Risk NEWS2: 7 or above Or significant clinical concerns Is this patient for Ward based care? Yes No Emergency Emergency Action! Action! Escalate to NIC Registered Nurse to review via Nervecentre. patient and Immediately medical notes. inform Speciality Reg If appropriate: via Nervecentre. Escalate to NIC Immediately via Nervecentre. bleep CCOT on Escalate to 888. Ward Doctor via Nervecentre. For immediate For immediate review ART review ART call 2222 call 2222 Outcome: Response time - immediate. Clearly document management plan, ceiling of care and review patients resuscitation status. Where appropriate failure to respond to treatment within 2 hrs after treatment commenced demands urgent discussion

with responsible consultant and where

appropriate referral to Critical Care

Observation Frequency:

Commence urine output monitoring

(bleep 269).

Minimum 30 minutes

NEWS2: 5 or 6 Is this patient for Ward based care? Urgent Action! Urgent Action! Registered Registered Nurse to review Nurse to review patient and patient and medical notes. medical notes. Escalate to NIC via Nervecentre. If appropriate: Escalate toWard · Escalate to NIC Doctor via via Nervecentre. Nervecentre. · Escalate toWard Escalate toCCOT Doctor via via Nervecentre. Nervecentre. Outcome: Response time 30 minutes. Review/ discussion must be documented in medical notes. Failure of NEWS2 to improve after 2 hours since treatment commenced must be escalated to Speciality Registrar or above and CCOT. Speciality Registrar or above is responsible for ensuring patient is discussed with the relevant consultant

and where appropriate Critical Care

Observation Frequency:

Commence urine output monitoring

Doctors (Bleep 269).

Minimum 1 hourly

Medium Risk

Low-Medium Risk NEWS2: Single parameter of 3 Urgent Action! Registered Nurse to review patient and medical notes. Escalate to NIC via Nervecentre. Escalate to Ward Doctor via Nervecentre. Outcome: Response time 30 minutes. Review by Ward Doctor who will further escalate if concerned and documented in medical notes. Re-escalate if patient not reviewed in 30minutes. Observation Frequency: Minimum 1 Hourly Commence urine output monitoring

Low Risk NEWS2: 0-4 If no clinical concerns continue routine observations If clinically concerned escalate to NIC/ Ward Doctor via Nervecentre Outcome if action required: Review by Ward Doctor if required and clear management plan documented in medical notes. Re-escalate If patient not reviewed by Ward Doctor within 1 hour. Observation Frequency: 0 Score = Minimum 12 hourly 1-2 Score = Minimum 6 hourly 3-4 Score = Minimum 4 hourly First 24 hours of admission = 4 hourly

17.00–09.00 Mon–Fri Including weekends and bank holidays

Sherwood Forest Hospitals
NHS Foundation Trust

* 08:00-09:00 bleep SHO3 via switch

Escalation of Observation Flowcharts - For King's Mill Site ONLY

High Risk

NEWS2: 7 or above Or significant clinical concerns

Is this patient for Ward based care?

Yes

Emergency Action!

 Registered Nurse to review patient and management plan

If appropriate:

- Registered Nurse to review patient and medical notes
- Escalate to NIC via Nervecentre
- Immediately inform HOOH/ on call Registrar*

For immediate review ART call 2222

Emergency Action!

Registered Nurse to review patient and management plan

If appropriate:

- NIC via Nervecentre
- Immediately inform HOOH/on call Registrar*
- Escalate via
 Nervecentre 24/7

For immediate review ART call 2222

Outcome:

- Response time immediate
- Clearly document management plan, ceiling of care and review patients resuscitation status
- Where appropriate failure to respond to treatment within 2 hrs after treatment commenced demands urgent discussion with responsible consultant and where appropriate referral to Critical Care (bleep 269)

Observation Frequency: Minimum 30 minutes Commence urine output monitoring

Medium Risk

NEWS2: 5 or 6

Is this patient for Ward based care?

Yes

Urgent Action!

 Registered Nurse to review patient and management plan

If appropriate:

- Escalate to NIC via Nervecentre
- Escalate to HOOH/SHO*

No

Urgent Action!

 Registered Nurse to review patient and management plan

If appropriate:

- Escalate to NIC via Nervecentre
- Escalate to HOOH/\$HO*
- Escalate via Nervecentre 24/7

Outcome:

- · Response time 30 minutes
- If no response and immediately concerned call ART on ext.2222
- Review/ discussion must be documented in medical notes
- Failure of NEWS2 to improve after 2 hours since treatment commenced must be escalated to Speciality Registrar or above and CCOT
- Speciality Registrar or above is responsible for ensuring patient is discussed with the relevant consultant and where appropriate Critical Care Doctors (Bleep 269)

Observation Frequency:

Minimum 1 hourly
Commence urine output monitoring

Low-Medium Risk

NEWS2:

Single parameter of 3

Urgent Action!

 Registered Nurse to review patient and management plan

If appropriate:

- Escalate to NIC via Nervecentre
- Escalate to HOOH/ SHO*

Outcome:

- · Response time 30 minutes
- Review by Ward Doctor who will further escalate if concerned and documented in medical notes
- Re-escalate if patient not reviewed in 30minutes

Observation Frequency:

Minimum 1 Hourly Commence urine output monitoring

Low Risk

NEWS2: 0-4

If no clinical concerns continue routine observations

OR

If clinically concerned escalate to NIC and HOOH/\$HO*as appropriate

Outcome if action required:

- Reviewed by Ward Doctor and clear management plan documented in medical notes
- Re-escalate If patient not reviewed by Ward Doctor within 1 hour

Observation Frequency:

0 Score = Minimum 12 hourly 1-2 Score = Minimum 6 hourly 3-4 Score = Minimum 4 hourly First 24 hours of admission = 4 hourly



Escalation of Observation Flowcharts – For Mansfield Site ONLY

High Risk

NEWS2: 7 or above Or significant clinical concerns

Emergency Action!

- Registered Nurse to urgently review patient
- Escalate to NIC via Nervecentre
- 08:00-17:00 (Mon-Fri) immediately phone the ward Doctor
- Out of hours including weekends and bank holidays immediately phone out of hours GP
- Contact Buddy Ward Call 9(999) for ambulance transfer to KMH

Outcome:

- Response time within 30 minutes
- Transfer to KMH where appropriate

Observation Frequency:

Minimum 30 minutes Commence urine output monitoring

Medium Risk

NEWS2: 5 or 6

Urgent Action!

- Registered Nurse to urgently review
- Escalate to NIC via Nervecentre
- 08:00-17:00 (Mon-Fri) inform the ward Doctor via Nervecentre
- Out of hours including weekends and bank holidays phone out of hours GP (9)999 ambulance call if appropriate for urgent transfer to KMH

Outcome:

- Response time 30 minutes
- Review/ discussion must be documented in medical notes
- Transfer to KMH if appropriate

Observation Frequency:

Minimum 1 hourly Commence urine output monitoring

Low-Medium Risk

NEWS2:

Single parameter of 3

Urgent Action!

- Registered Nurse to review patient
- Escalate to NIC via Nervecentre
- 08:00-17:00 (Mon-Fri) inform the wards Doctor via Nervecentre
- Out of hours including weekends and bank holidays phone out of hours GP

Outcome:

- Response time 30 minutes
- Review by Doctor/ACP who will further escalate if concerned and document in medical notes
- Re-escalate if patient not reviewed in 30minutes

Observation Frequency:

Minimum 1 Hourly Commence urine output monitoring

Low Risk

NEWS2: 0-4

Is Action required?

Score 0 - Continue with routine observations

Score 1-4 - Registered nurse to review patient and consider if increased frequency of monitoring and/ or escalation of patient required

Outcome if action required:

- 08.00-17.00 (Mon-Fri) Phone ward
- Out of hours including weekends and bank holidays phone out of hours GP

Observation Frequency:

0 Score = Minimum 24 hourly 1-2 Score = Minimum 6 hourly

3-4 Score = Minimum 4 hourly



Escalation of Observation Flowcharts – For Newark Site ONLY

High Risk

NEWS2: 7 or above Or significant clinical concerns

Emergency Action!

- Registered Nurse to urgently review patient
- · Escalate to NIC via Nervecentre
- 08:00-19:00 (Mon-Fri) immediately bleep ward doctor
- Out of hours including weekends and bank holidays immediately call UCC doctor

Call 9(999) for EMAS ambulance transfer to KMH. If EMAS can't respond call Ambicorp (01623 880988) for an urgent transfer

Cardiac Arrest: Call 2222

Outcome:

- Response time within 30 minutes
- Transfer to KMH where appropriate

Observation Frequency: Minimum 30 minutes Commence urine output monitoring

Medium Risk

NEWS2: 5 or 6

Urgent Action!

- Registered Nurse to urgently review patient
- Escalate to NIC via Nervecentre
- 08:00-19:00 (Mon-Fri) inform ward doctor through Nervecentre
- Out of hours including weekends and bank holidays call UCC doctor
- Consider early transfer to KMH site through discussion with intensivist on ICU (Contactable through switchboard)

Outcome:

- Response time 30 minutes
- Review/ discussion must be documented in medical notes
- · Transfer to KMH if appropriate

Observation Frequency:

Minimum 1 hourly Commence urine output monitoring

Low-Medium Risk

NEWS2:

Single parameter of 3

Urgent Action!

- Registered Nurse to review patient
- · Escalate to NIC via Nervecentre
- 08:00-19:00 (Mon-Fri) inform ward doctor via Nervecentre
- Out of hours including weekends and bank holidays call UCC doctor

Outcome:

- Response time 30 minutes
- Reviewed by Doctor and clear management plan documented in medical notes
- Re-escalate if not reviewed/ discussed in 30 minutes

Observation Frequency:

Minimum 1 Hourly
Commence urine output monitoring

Low Risk

NEWS2: 0-4

Is Action required?

Score 0 – Continue with routine observations

Score 1-4 – Registered nurse to review patient and consider if increased frequency of monitoring and/ or escalation of patient required

Outcome if action required:

- 08:00-19:00 (Mon-Fri) inform ward doctor
- Out of hours including weekends and bank holidays call UCC doctor

Observation Frequency:

0 Score = Minimum 24 hourly 1-2 Score = Minimum 6 hourly

3-4 Score = Minimum 4 hourly

000



Escalation of Observation Flowcharts – For The Emergency Department ONLY

High Risk

NEWS2: 7 or above Or significant clinical concerns

Emergency Action!

- Registered Nurse to review patient and medical notes
- Escalate to NIC
- Escalate to responsible clinician, consultant in charge or middle grade or above in charge overnight Transfer to Resus

.....

Outcome:

- Response time within 30 minutes
- Discussion with consultant/ middle grade or above
- Clearly document management plan, ceiling of care and review patients resuscitation status
- Consider referral to speciality registrar and on call ED anaesthetist at earliest opportunity

Observation Frequency: Minimum 30 minutes Commence urine output monitoring

Medium Risk

NEWS2: 5 or 6

Urgent Action!

- Registered Nurse to review patient and medical notes
- Escalate to NIC
- Escalate to responsible clinician, consultant in charge or middle grade or above in charge overnight

Consider transferring patient to Resus

Outcome:

- Response time 30 minutes
- Review/ discussion must be documented
- Failure of NEWS2 score to improve after treatment commenced must be escalated to consultant in charge/ middle grade or above in charge if they are not the responsible clinician
- Consider presenting problem; increased frequency of observations may be required due to trauma, interventions or rapidly changing conditions

Observation Frequency:

Minimum 30 minutes
Commence urine output monitoring

Low-Medium Risk

NEWS2:

Single parameter of 3

Urgent Action!

- Registered Nurse to review medical plan
- Escalate to NIC
- Escalate to responsible ED clinician

Outcome:

- Response time 30 minutes
- Review by Doctor/ACP who will further escalate if concerned and document in medical notes
- Re-escalate if patient not reviewed in 30minutes

Observation Frequency: Minimum 1 Hourly Commence urine output monitoring

Low Risk

NEWS2: 0-4

If no clinical concerns continue routine observations

If clinically concerned escalate to:

- Registered nurse who will review patient and medical plan
- NIC
- Responsible clinician, consultant in charge or middle grade or above in charge overnight

Outcome if action required:

- Reviewed by Doctor/ACP if required and clear management plan documented in medical notes
- Re-escalate If patient not reviewed within 1 hour
- Consider presenting condition as more frequent observations may be required

Observation Frequency:

0 Score = Observations to be repeated prior to discharge/ transfer to ward or if any change in condition

1-4 Score = Observations to be repeated within the first 2hrs of arrival and before discharge/ transfer to ward or if change in condition

Appendix 5

NEWS2 Score:

Physiological	Score						
parameter	3	2	1	0	1	2	3
Respiration rate (per minute)	≤8		9–11	12–20		21–24	≥25
SpO ₂ Scale 1 (%)	≤91	92–93	94-95	≤96			
SpO ₂ Scale 2 (%)	≤83	84–85	86–87	88–92 ≤93 on air	93–94 on oxygen	95–96 on oxygen	≥97 on oxygen
Air or oxygen?		Oxygen		Air			
Systolic blood pressure (mmHg)	≤90	91–100	101-110	111–219			≥220
Pulse (per minute)	≤40		41–50	51–90	91–110	111–130	≥131
Consciousness				Alert			CVPU
Temperature (°C)	≤35.0		35.1–36.0	36.1–38.0	38.1–39.0	≥39.1	

Appendix 6

Post-operative patient monitoring for Adult Surgical In-Patients:

This post-operative patient monitoring procedure directs the measurement of patient vital signs for the first six hours on return to the ward after surgery. All other areas including Day Case and Endoscopy should follow Day Case Surgical Pathways. For General Surgery patients please also refer to: Nursing Care Pathway for Patients Undergoing General Surgery Procedures.

- 1. Immediately on return to the ward: Complete a full set of mandatory vital signs:
 - Respiratory rate (RR)
 - Oxygen saturations (SpO2)
 - Inspired O2
 - Blood pressure (BP)
 - Heart rate (HR)
 - Urine output
 - Level of Consciousness using ACVPU
 - Temperature (T)
 - Pain score
 - **Bowels**

Compare this initial set of vital signs with those performed pre-operations and in recovery and check for improvement or deterioration.

Repeat these initial observations every 30 minutes for 2 hours and escalate any concerns according to the Observations and Escalation Policy.

In addition to vital signs, carry out surgery-specific observations including:

- Wound site
- Wound drainage and record on fluid balance chart
- Urine output and record on fluid balance chart
- IV fluid administration
- Blood glucose levels
- Nausea
- 2. After 2 hours of 30 minute observations* (if the cumulative NEWS2 is ≤4and there is no single parameter score of 3) reduce the frequency of observations to hourly. Escalate according to the Observations and Escalation Policy.

- *Day surgery patients may have observations discontinued sooner if all discharge criteria have been achieved - refer to Day Surgery guidance.
- 3. After 4 hours of hourly observations (if the cumulative NEWS2 is ≤4 and there is no single parameter score of 3) reduce observations further to be in line with the Observations and Escalation Policy. From this point onwards the frequency of observations is guided by the NEWS2 score.

Please note: The NEWS2 score is only one indicator of the patient's physiological status. Please make a comprehensive assessment of the patient throughout their recovery period and, if in doubt, escalate any patient you are worried about.

Version: 9.0; Issued: June 2023 Page 30 of 34

Appendix 7

SBAR Communication Tool:

S

Situation:

I am (name), (X) nurse on ward (X)
I am calling about (patient X)
I am calling because I am concerned that...
(e.g. BP is low/high, pulse is XX, temperature is XX, Early Warning Score is XX)

B

Background:

Patient (X) was admitted on (XX date) with... (e.g. Ml/chest infection)

They have had (X operation/procedure/investign

They have had (X operation/procedure/investigation)
Patient (X)'s condition has changed in the last (XX mins)
Their last set of obs were (XX)

Patient (X)'s normal condition is... (e.g. alert/drowsy/confused, pain free)

A

Assessment:

I think the problem is (XXX)

And I have...

(e.g. given O₂/analgesia, stopped the infusion)

I am not sure what the problem is but patient (X) is deteriorating

OR

I don't know what's wrong but I am really worried

R

Recommendation:

I need you to...

Come to see the patient in the next (XX mins)

Is there anything I need to do in the mean time? (e.g. stop the fluid/repeat the obs)

Ask receiver to repeat key information to ensure understanding

Title: Observations and Escalation Policy for Adult Patients
Version: 9.0; Issued: June 2023 Page **31** of **34**



APPENDIX 8 - EQUALITY IMPACT ASSESSMENT FORM (EQIA)

Name of service/police	cy/procedure being reviewed: Observation	s and Escalation Policy for Adult Patients		
New or existing servi	ce/policy/procedure: Existing			
Date of Assessment:	28 th April 2023			
	y/procedure and its implementation answ r implementation down into areas)	ver the questions a - c below against each characte	eristic (if relevant consider	
a) Using data and supporting information, what issues, needs or Characteristic barriers could the protected characteristic groups' experience? For example, are there any known health inequality or access issues to consider?		b) What is already in place in the policy or its implementation to address any inequalities or barriers to access including under representation at clinics, screening?	c) Please state any barriers that still need to be addressed and any proposed actions to eliminate inequality	
The area of policy or	its implementation being assessed:			
Race and Ethnicity	None	Clear guidelines in terms of implementing the policy	None	
Gender	None	Clear guidelines in terms of implementing the policy	None	
Age	Paediatric and neonatal Patients are excluded from this policy.	See PEWS Escalation.	None	
Religion	None	Clear guidelines in terms of implementing the policy	None	
Disability	None	Clear guidelines in terms of implementing the policy	None	
Sexuality	None	Clear guidelines in terms of implementing the policy	None	
Pregnancy and Maternity	Maternity patients are excluded from this policy.	See Maternity Early Warning Scoring (MEWS) Guideline.	None	
Gender Reassignment	None	Clear guidelines in terms of implementing the policy	None	
Marriage and Civil Partnership	None	Clear guidelines in terms of implementing the policy	None	
Socio-Economic Factors (i.e. living in a poorer neighbourhood / social deprivation)	None	Clear guidelines in terms of implementing the policy	None	

Title: Observations and Escalation Policy for Adult Patients Version: 9.0; Issued: June 2023

Page **32** of **34**



What consultation with protected characteristic groups including patient groups have you carried out?

• End user staff groups, Nursing and Midwifery Board, Specialist Teams and Deteriorating Patient Group.

What data or information did you use in support of this EqIA?

· Reviewed in line with national guidance

As far as you are aware are there any Human Rights issues be taken into account such as arising from surveys, questionnaires, comments, concerns, complaints or compliments?

• No

Level of impact

From the information provided above and following EQIA guidance document Guidance on how to complete an EIA (<u>click here</u>), please indicate the perceived level of impact:

Low Level of Impact

For high or medium levels of impact, please forward a copy of this form to the HR Secretaries for inclusion at the next Diversity and Inclusivity meeting.

Name of Responsible Person undertaking this assessment

Alison Davidson

Signature:

A.Davidson

Date:

28th April 2023

Title: Observations and Escalation Policy for Adult Patients
Version: 9.0; Issued: June 2023 Page **33** of **34**



APPENDIX 9 – ENVIRONMENTAL IMPACT ASSESSMENT

The purpose of an environmental impact assessment is to identify the environmental impact, assess the significance of the consequences and, if required, reduce and mitigate the effect by either, a) amend the policy b) implement mitigating actions.

Area of impact	Environmental Risk/Impacts to consider	Yes/No	Action Taken (where necessary)
Waste and materials	 Is the policy encouraging using more materials/supplies? Is the policy likely to increase the waste produced? Does the policy fail to utilise opportunities for introduction/replacement of materials that can be recycled? 	No No No	NA
Soil/Land	 Is the policy likely to promote the use of substances dangerous to the land if released? (e.g. lubricants, liquid chemicals) Does the policy fail to consider the need to provide adequate containment for these substances? (For example bunded containers, etc.) 	No No	NA
Water	 Is the policy likely to result in an increase of water usage? (estimate quantities) Is the policy likely to result in water being polluted? (e.g. dangerous chemicals being introduced in the water) Does the policy fail to include a mitigating procedure? (e.g. modify procedure to prevent water from being polluted; polluted water containment for adequate disposal) 	No No	NA
Air	 Is the policy likely to result in the introduction of procedures and equipment with resulting emissions to air? (For example use of a furnaces; combustion of fuels, emission or particles to the atmosphere, etc.) Does the policy fail to include a procedure to mitigate the effects? Does the policy fail to require compliance with the limits of emission imposed by the relevant regulations? 	No No No	NA
Energy	Does the policy result in an increase in energy consumption levels in the Trust? (estimate quantities)	No	NA
Nuisances	 Would the policy result in the creation of nuisances such as noise or odour (for staff, patients, visitors, neighbours and other relevant stakeholders)? 	No	NA